

AD-A131 086

THE POSSIBILITY OF AMPLIFYING INFRARED RADIATION BY
HIGH-PRESSURE REACTING GAS(U) FOREIGN TECHNOLOGY DIV
WRIGHT-PATTERSON AFB OH V A KOCHLAN ET AL. 22 JUN 83
FTD-ID(RS)T-0569-83

1/1


UNCLASSIFIED

F/G 17/5

NL



END
DATE
FILMED
#3 H.S
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 131 086



FOREIGN TECHNOLOGY DIVISION



THE POSSIBILITY OF AMPLIFYING INFRARED RADIATION BY
HIGH-PRESSURE REACTING GAS

by

V.A. Kochelan, Yu.A. Kukibnyy



DTIC FILE COPY

Approved for public release;
distribution unlimited.

88 08 03 049

EDITED TRANSLATION

FTD-ID(RS)T-0569-83

22 June 1983

MICROFICHE NR: FTD-83-C-000767

THE POSSIBILITY OF AMPLIFYING INFRARED RADIATION BY
HIGH-PRESSURE REACTING GAS

By: V.A. Kochelan, Yu.A. Kukibnyy

English pages: 7

Source: Ukrainskiy Fizicheskiy Zhurnal, Vol. 17,
Nr. 1, January 1972, pp. 149-151

Country of origin: USSR

Translated by: Robert D. Hill

Requester: FTD/TQTD

Approved for public release; distribution unlimited.



THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP.AFB, OHIO.

FTD -ID(RS)T-0569-83

Date 22 Jun 19 83

U. S. BOARD ON GEOGRAPHIC NAMES transliteration SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
А а	<i>А а</i>	A, a	Р р	<i>Р р</i>	R, r
Б б	<i>Б б</i>	B, b	С с	<i>С с</i>	S, s
В в	<i>В в</i>	V, v	Т т	<i>Т т</i>	T, t
Г г	<i>Г г</i>	G, g	У у	<i>У у</i>	U, u
Д д	<i>Д д</i>	D, d	Ф ф	<i>Ф ф</i>	F, f
Е е	<i>Е е</i>	Ye, ye; E, e*	Х х	<i>Х х</i>	Kh, kh
Ж ж	<i>Ж ж</i>	Zh, zh	Ц ц	<i>Ц ц</i>	Ts, ts
З э	<i>З э</i>	Z, z	Ч ч	<i>Ч ч</i>	Ch, ch
И и	<i>И и</i>	I, i	Ш ш	<i>Ш ш</i>	Sh, sh
Й й	<i>Й й</i>	Y, y	Щ щ	<i>Щ щ</i>	Shch, shch
К к	<i>К к</i>	K, k	Ъ ъ	<i>Ъ ъ</i>	"
Л л	<i>Л л</i>	L, l	Ы ы	<i>Ы ы</i>	Y, y
М м	<i>М м</i>	M, m	Ь ь	<i>Ь ь</i>	'
Н н	<i>Н н</i>	N, n	Э э	<i>Э э</i>	E, e
О о	<i>О о</i>	O, o	Ю ю	<i>Ю ю</i>	Yu, yu
П п	<i>П п</i>	P, p	Я я	<i>Я я</i>	Ya, ya

*ye initially, after vowels, and after Ъ, ь; e elsewhere.
When written as ё in Russian, transliterate as yë or ë.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh ⁻¹
cos	cos	ch	cosh	arc ch	cosh ⁻¹
tg	tan	th	tanh	arc th	tanh ⁻¹
ctg	cot	cth	coth	arc cth	coth ⁻¹
sec	sec	sch	sech	arc sch	sech ⁻¹
cosec	csc	csch	csch	arc csch	csch ⁻¹

Russian English

rot curl
lg log

GRAPHICS DISCLAIMER

All figures, graphics, tables, equations, etc. merged into this translation were extracted from the best quality copy available.

THE POSSIBILITY OF AMPLIFYING INFRARED RADIATION BY HIGH-PRESSURE
REACTING GAS

V.A. Kochelan, Yu.A. Kukibnyy

Being intensely investigated at present are infrared chemical lasers in which the process of the stimulated light emission is preceded by a chemical reaction, as a result of which populated molecules are inversely formed [1]. Usually used in such type of lasers are reacting gases of low pressure, since with an increase in pressure the rate of deactivation of the excited molecules is increased.

Noted in this work is the possibility of amplifying the infrared radiation by a reacting gas in cases when serving as the working transition is the phototransition of a pair of reacting molecules, which lead to a change in their translational-rotational motion with an unchanged electron state. It is found that in such cases reacting gases of high pressure can be used for the light amplification. This is the special feature of the considered mechanism of light

amplification and a number of others are analogous to properties of the chemical laser on electron phototransitions proposed in work [2].




Figure.

For definiteness, let us examine the reaction of the recombination of atoms of two kinds. A pair of reacting particles, in approaching, moves in the effective field of the electrons and nuclei $U_k(r) = U_0(r) + \frac{\hbar^2 k(k+1)}{2\mu r^2}$ with an energy of the relative motion $E > 0$ (k - rotational quantum number, μ - normalized mass, see the figure). At reciprocal distances $r \approx r_{0k}$ there appears the dipole moment $p(r)$, and the phototransition becomes possible. In the case of collisions with $E < \hbar\omega$ (ω - light frequency), the emission of a quantum leads to the formation of a stable molecule with energy $E_{\text{mol}} = E - \hbar\omega < 0$, v - the oscillating quantum number. This phototransition is more probable than the absorption $E \rightarrow E + \hbar\omega$ ($E > 0$) and radiation $E \rightarrow E - \hbar\omega$ (when $E > \hbar\omega$), since for it the final states of the transition are characterized by localized wave functions. Considering the

distribution of atoms with respect to energies, we can be sure that in a gas with temperature T the emission processes predominate over the absorption processes when $\hbar\omega > T$ (T - temperature in energy units). Below we will assume that this condition is fulfilled. If, moreover, the rapid relaxation of the excited levels E_{nk} is ensured, then the light amplification factor in the recombining gas is $\alpha > 0$. In cases when we can disregard the absorption by bound molecules, where $\alpha = a n_1 n_2 / n_1$, and n_1 and n_2 are concentrations of the recombining atoms. Coefficient a is usually [3] expressed in terms of matrix elements $\rho_{12}(\omega) = \langle E_{nk}, k \pm 1 | \rho(r) | E, k \rangle$. The latter, for transitions between the quasi-classical states, depend on the difference $E - E_{nk} = \hbar\omega$ [4]; the dependence of k on E_1, E_2 separately is insignificant. Let us give the expression for a in the following cases. When $T \ll \hbar\omega_{nk} = E_{n+1,k} - E_{nk}$ the phototransitions should be observed mainly on rotational sublevels of one oscillating state $E_{nk} \approx E_{n0} + B_r k(k+1)$, for which $E_{n0} \approx \hbar\omega$, then

$$a \approx \frac{(2\pi)^{7/2} \hbar^4 \omega_{nk} \omega}{3 c \mu^{3/2} T^{1/2} B_r} |\rho_{12}(\omega)|^2. \quad (1)$$

In the case of $T > \hbar\omega_{nk}$ phototransitions are observed on the many oscillating levels, and the expression for a takes the form

$$a \approx \frac{(2\pi)^{7/2} \hbar^4 \omega_{nk} \omega}{6 c (\mu T)^{3/2}} \sigma(T) |\rho_{12}(\omega)|^2. \quad (2)$$

Factor $\sigma(T)$ appears due to the presence of the centrifugal barrier in the potential energy $U_k(r)$ [5] and is determined by the equality

$$\sigma(T) = \sum_k (2k+1) \exp \left[-\frac{U_k(r_k^*)}{T} \right].$$

r_k^* - the position of the maximum of the barrier (see the figure).

The matrix element $p_{12}(\omega)$ can be evaluated, having used the Morse model:

$$U_0(r) = U_0 [e^{-2\beta(r-r_0)} - 2e^{-\beta(r-r_0)}].$$

For the most urgent case $\hbar\omega \ll U_0$ the expression for $p_{12}(\omega)$ can be found from the formula of work [6]:

$$|p_{12}(\omega)|^2 \approx \left(\frac{dp}{dr} \Big|_{r_0} \right)^2 \frac{1}{\beta^2 \hbar^2 \omega^2} e^{-\frac{2\omega}{\omega_0}} \quad (3)$$

where $\omega_0 = \sqrt{\frac{\beta^2 U_0}{2\mu}}$ is the frequency of oscillations of the main state. Hence it follows that the most probable is the phototransition at frequencies $\omega \leq \omega_0$. In the same model it is possible to find $\hbar\omega_{00} \approx 2 \sqrt{\frac{\beta^2 \hbar^2 \omega}{2\mu}}$.

If in the working volume there are reaction products, then the light amplification factor is written in the form

$$\alpha = \alpha n_1 n_2 \xi, \quad \xi = 1 - \frac{\delta m}{n_1 n_2} \left(\frac{\mu T}{2\pi \hbar^2} \right)^{3/2}, \quad (4)$$

where $\delta = T/B_0$ when

$T > \hbar\omega_{00}$ and $\delta = 1/\sigma(T)$ when $T < \hbar\omega_{00}$; m - concentration of the molecules - reaction products in states with energies $|E_{00}| \leq \hbar\omega$. In conditions of the recombining gas, these states are populated with the speed of the thermal recombination reaction $kn_1 n_2$. If their destruction is ensured with a rate of gm molecules per unit time, then the condition of amplification $\xi > 0$ takes the form

$$\frac{g}{k} > \delta \left(\frac{\mu T}{2\pi \hbar^2} \right)^{3/2}. \quad (5)$$

Under a pressure of the order of atmospheric, a sufficient rate of deactivation of the nonequilibrium populated levels with $|E_{00}| \leq \hbar\omega$ can be ensured by the inelastic collisions in the gas,

since for these levels $\hbar\omega_{0k} \ll T$, the deactivation occurs as a result of one or several gas-kinetic collisions. It is essential that here with the growth in the pressure, the necessary condition of light amplification (5) is not deteriorated, since g and k are proportional to the total concentration of particles in the gas N ; at the same time $\alpha \sim N^2$. Consequently, the effectiveness of the examined light amplification mechanism is increased with an increase in pressure.

Let us estimate a and α for the specific recombination reaction $\text{Li} + \text{H} \rightarrow \text{LiH} + \hbar\omega$. For the molecule LiH , it is possible to assume that $\hbar\omega_0 = 1400 \text{ cm}^{-1}$, $\left(\frac{dp}{dr}\right)_e = 0.3e$, e - the electron charge [7]. Assuming that $T = 500^\circ\text{K}$ and $\omega = \omega_0/2$, we can find that $\hbar\omega_{0k} < T$, so that it follows to use formula (2). Let us estimate the value of σ by assuming that $U_e(r)$ has the form of the Morse potential with parameters of the basic electron state LiH , and, as result, we get $\sigma = 1.1 \cdot 10^{-3}$, $a = 2.5 \times 10^{-11} \text{ cm}^3$. For g we take the ^{value} A determined by the cross section of the usual gas-kinetic collision $\frac{g}{N} = 10^{-10} \text{ s}^{-1}$; then from (5) there follows the condition which must be satisfied by the rate of the thermal recombination reaction: $k < N \cdot 10^{-11} \text{ cm}^3 \cdot \text{s}^{-1}$. This inequality is apparently fulfilled, since usually for the three-particle recombination reactions $k/N \ll 10^{-11} - 10^{-12} \text{ cm}^3 \cdot \text{s}^{-1}$. Assuming that $\xi = 1$; $n_1 = n_2 = 10^{11} \text{ cm}^{-3}$, for the light amplification factor we get $\alpha = 2.5 \cdot 10^{-3} \text{ cm}^{-1}$. It is evident that the light amplification factor is sufficient for carrying out the laser generation.

The authors express their deep appreciation to S.I. Pekar for his valuable remarks made during the reading of the manuscript.

Institute of Semiconductors of the Academy of Sciences of the USSR,
Kiev - - Received 29 June 1971

References

1. N.G. Basov, V.I. Igoshin, Ye.P. Markin, A.N. Orayevskiy.
Collection: "Kvantovaya elektronika" [Quantum Electronics], Edited by
N.G. Basov, No. 2, 1971, page 3.
2. S.I. Pekar, DAN [Reports of Academy of Sciences] of USSR, 187,
555, 1969.
3. V.A. Kochelap, UFZh [Ukrainian Physics Journal], 15, 1001; 1213,
1970.
4. L.D. Landau, Ye.M. Lifshits, "Kvantovaya mekhanika" [Quantum
Mechanics], Moscow, Fizmatgiz, 1963.
5. V.A. Kochelap, S.I. Pekar. ZhETF [Journal of Experimental and

Technical Physics], 58, 854, 1970; UFZh, 15, 1057, 1970.

6. G.A. Askar'yan. ZhETF, 48, 667, 1965.

7. Ch.F. Bender, E.R. Davidson. J. Chem. Phys., 49, 4222, 1968.

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

<u>ORGANIZATION</u>	<u>MICROFICHE</u>
A205 DMAHTC	1
A210 DMAAC	1
B344 DIA/RTS-2C	9
C043 USAMIIA	1
C500 TRADOC	1
C509 BALLISTIC RES LAB	1
C510 R&T LABS/AVRADCOM	1
C513 ARRADCOM	1
C535 AVRADCOM/TSARCOM	1
C539 TRASANA	1
C591 FSTC	4
C619 MIA REDSTONE	1
D008 NISC	1
E053 HQ USAF/INET	1
E403 AFSC/INA	1
E404 AEDC/DOF	1
E408 AFWL	1
E410 AD/IND	1
E429 SD/IND	1
P005 DOE/ISA/DDI	1
P050 CIA/OCR/ADD/SD	2
AFIT/LDE	1
FTD	
CCN	1
NIA/PHS	1
NIIS	2
LLNL/Code L-389	1
NASA/NST-44	1
NSA/1213/TDL	2

END

DATE
FILMED

9 - 83

DTIC